

J. Linn. Soc. Bot. 10:183. 1868.

ACROPORIUM, gen. nov.

Caulis primarius repens, ramos elongatos ramosos in caespites congestos prodens. Folia nitida, undique inserta, uniformia, in ramorum apicibus cuspidatim imbricata, cellulis elongatis alaribus magnis conspicuis. Fructus e partibus superioribus ramorum oriens. Theca in pedunculo gracillimo, minuta, ovalis, inclinata, operculo longirostro, peristomio *Stereodontis*.

A. BREVICUSPIDATUM, sp. n. Monoicum; caulis inaequaliter pinnatim ramosus, elongatus; folia patentia laxè imbricata, in ramorum apicibus in cuspidem brevem congesta, late oblongo-ovalia, breviter acuminata, concava, margine versus apicem incurva, integerrima, cellulis angustis laevibus pellucidis, alaribus conspicuis aurantiacis; perichæetalia parva, caulinis conformia, integerrima; theca in pedunculo elongato gracillimo apice scabro, oblonga, parva, inclinata, peristomio interno ciliis singulis processibus brevioribus?

Hab. Manua, on large trees (1500 ft.). No. 137. Also from the Sandwich Islands and South-Sea Islands, *herb. Hooker*.

Corresponding in size, habit, and appearance with *A. turgidum*, Dozy et Molk. (*Hypnum*); inflorescence, however, monoecious, and foliage paler and softer.