

TERMS OF USE

This pdf is provided by Magnolia Press for private/research use.
Commercial sale or deposition in a public library or website is prohibited.



Zootaxa 2992: 61–68 (2011)
www.mapress.com/zootaxa/
Copyright © 2011 · Magnolia Press

Article

ISSN 1175-5326 (print edition)

ZOOTAXA

ISSN 1175-5334 (online edition)

Diplocentrus bicolor sp. n. (Scorpiones: Diplocentridae) from Jalisco, Mexico

GERARDO A. CONTRERAS-FÉLIX¹ & CARLOS E. SANTIBÁÑEZ-LÓPEZ²

Colección Nacional de Arácnidos. Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México;
Apartado Postal 70-153, México D.F. 04510. MEXICO. E-mail: ¹contrerasfelixga@hotmail.com; ²ironc81@hotmail.com

Abstract

Diplocentrus bicolor sp. n. from Huejuquilla el Alto, in northern Jalisco and two nearby localities in Zacatecas is described. It is compared to its geographically closer species from Zacatecas, Aguascalientes and Nayarit. A map with its known distribution is provided.

Key words: scorpion, Diplocentridae, *Diplocentrus bicolor*, new species, Mexico, Jalisco

Introduction

The genus *Diplocentrus* Peters, 1861 holds the highest diversity within the family Diplocentridae with 51 species (Santibáñez-López & Francke, 2008; Santibáñez-López *et al.*, 2011). Its distribution ranges from the southern states of the USA to northern Honduras (Sissom & Fet, 2000; Francke & Ponce-Saavedra, 2005). It is, however, poorly sampled in the Mexican state of Jalisco where only one species had been reported previously near the border with Nayarit (*D. gertschi* Sissom & Walker, 1992). Despite the recent taxonomic work on the genus (Francke & Ponce-Saavedra, 2005; Francke, 2007; Santibáñez-López & Francke, 2008; Francke & Quijano-Ravell, 2009; Santibáñez-López *et al.*, 2011) the knowledge of the genus in Mexico is incomplete. The present contribution provides the description of the second species of the genus *Diplocentrus* known from Jalisco and adjacent southern Zacatecas.

Material and methods

Nomenclature and mensuration follows Stahnke (1970), except for trichobothrial terminology after Vachon (1974), metasomal and pedipalpal carinal terminology after Francke (1977). Surfaces of the pedipalp, carapace, mesosoma and metasoma were observed as in Santibáñez-López & Sissom (2010). Higher level taxonomy of scorpions follows Coddington *et al.* (2004) and Prendini & Wheeler (2005). Photography of the female and male carapace, pedipalp femur, patella and chela under black light follows Prendini (2003) and Volschenk (2005). Measurements were taken with an ocular micrometer calibrated at 10X and are given in millimeters. Abbreviations for depositaries: AMNH—American Museum of Natural History, New York. CNAN—Colección Nacional de Arácnidos, Instituto de Biología, Universidad Nacional Autónoma de México, México D.F. México.

TERMS OF USE

This pdf is provided by Magnolia Press for private/research use.
Commercial sale or deposition in a public library or website is prohibited.

Taxonomy**Family DIPLOCENTRIDAE****Genus *Diplocentrus******Diplocentrus bicolor* sp. n.**

(Figures 1–10)

Etymology. The specific epithet refers to the contrasting coloration of the darker body with the legs considerably lighter in color; and it is used as a noun in apposition.

Types. Jalisco: Municipio Huejuquilla el Alto: El Vallecito 15 km from Huejuquilla el Alto along the road to San Juan Capistrano (N 22° 40.27', W 103° 57.63', 1233 m) 6.vii.2005. O. Francke, J. Ponce, M. Córdova, A. Jaimes, G. Francke & V. Capovilla. 1 ♂ (CNAN-T0683).

Paratypes: **Jalisco:** Municipio Huejuquilla el Alto: El Vallecito 15 km from Huejuquilla el Alto along the road to San Juan Capistrano (N 22° 40.27', W 103° 57.63', 1233 m) 6.vii.2005. O. Francke, J. Ponce, M. Córdova, A. Jaimes, G. Francke & V. Capovilla. 1 ♂, 2 ♀ (CNAN-T0684). **Jalisco:** Municipio Mezquitic: Carretera a Mezquitic between Mezquitic and Ojuelos (N 22° 21.601', W 103° 38.622', 2089 m) 7.vii.2005. O. Francke, J. Ponce, M. Córdova, A. Jaimes, G. Francke & V. Capovilla. 2 ♂ (AMNH). **Jalisco:** Municipio Huejuquilla el Alto: El Vallecito 18 km from Huejuquilla el Alto along road to San Juan Capistrano (N 22° 40.277', W 103° 57.638' 1233 m) 6.vii.2005. . O. Francke, J. Ponce, M. Córdova, A. Jaimes, G. Francke & V. Capovilla. 1 ♀ (CNAN-T0684).

Other specimens examined: **Jalisco:** Municipio Huejuquilla el Alto: El Vallecito 15 km from Huejuquilla el alto along the road to San Juan Capistrano (N 22° 40.27', W 103° 57.63', 1233 m) 6.vii.2005. O. Francke, J. Ponce, M. Córdova, A. Jaimes, G. Francke & V. Capovilla. 4 ♂ (CNAN-S03083); 3 ♂ (AMNH). **Jalisco:** Municipio Mezquitic: Carretera a Mezquitic between Mezquitic and Ojuelos (N 22° 21.601', W 103° 38.62' 2089 m) 7.vii.2005. O. Francke, J. Ponce, M. Córdova, A. Jaimes, G. Francke & V. Capovilla. 4 ♂ (CNAN-S03049). 3 ♀ (CNAN-S03050) 5 ♂ and 4 ♀ (AMNH). **Jalisco:** Municipio Mezquitic: 4 km to North from Mezquitic (N 22° 24.732', W 103° 43.222', 1388 m) 7.vii.2005 . O. Francke, J. Ponce, M. Córdova, A. Jaimes, G. Francke & V. Capovilla. 2 ♂ (CNAN-S03084) 1 ♂ 1 ♀ (AMNH). **Zacatecas:** Municipio Valparaíso: 4.1 km north from San Juan Capistrano (N 22° 40' 44", W 104° 06' 22", 1169 m) without date. Unknown colector. 1 ♂ (CNAN-S03029). **Zacatecas:** Municipio Valparaíso: Deviation to Los Tanques (N 22° 40' 14", W 104° 02' 32", 1099 m) 19.ix.2001. E. Gonzalez. 4 ♂, 1 ♀ (CNAN-S03030).

Distribution. Known from two localities, 3 km from each other, in northern Jalisco and from another two in southern Zacatecas (Fig. 11).

Diagnosis. Adults 60 to 70 mm long. Brownish to dark red, legs pale yellow to cream, contrasting sharply with the rest of the body. Frontal notch in anterior margin of carapace "U" shaped, notch moderately deep shagreened. Pedipalp femur orthobothrioxic, wider than deep, dorsal surface flat, weakly to sparsely granulose. Pedipalp patella orthobothrioxic; dorsal external carina weak to moderate, smooth in males; ventral submedian carina weak to faint, smooth. Chela orthobothrioxic; digital carina strong, smooth ending at the base of the fixed finger in males; dorsal surface slightly reticulate to smooth; pedipalp carination weaker and smoother in females. Telotarsal formula 5–6/6–7: 6/7: 7/8: 7/8. Pectinal tooth count on males (n=29) 17–19 (mode= 18); on females (n=10) 14–16 (mode= 15).

Diplocentrus bicolor sp. n. is similar to *D. poncei* Francke & Quijano-Revel, 2009 in overall size; the frontal notch of the carapace is "U" shaped in both species; the femur is wider than deep in both species and the dorsal surface flat, and both are similar in pectinal tooth counts. It is geographically closest westward to *D. gertschi* from Nayarit (Sissom & Walker, 1992), sharing a similar telotarsal formula (on the last leg 7/8), the dorsal surface of pedipalp wider than deep, and the pedipalp patella dorsal external carina weak to moderate and smooth in males; it is also closest geographically to the north and the east to *D. zacatecanus* Hoffmann, 1932, from Zacatecas, San Luis Potosí, Querétaro, Michoacán, Hidalgo, Guanajuato, Estado de México, Durango y Aguascalientes (Ponce-Saavedra *et al.*, 2009) sharing similar femur dorsal surface shape and similar carination development in the pedipalp patella. *D. bicolor* sp. n. can be clearly distinguished from *D. poncei* by the orthobothrioxic condition on the pedipalp patella and chela (*D. poncei* is neobothrioxic on both); by a higher telotarsal formula (5/7 5/7 on the last two legs in *D. poncei* whereas on *D. bicolor* sp.n. they are 7/8:7/8). Digital carina in the pedipalp chela in males

TERMS OF USE

This pdf is provided by Magnolia Press for private/research use.
Commercial sale or deposition in a public library or website is prohibited.



FIGURES 1–2. Habitus of holotype male. 1. Dorsal view. 2. Ventral view. Scale bar= 10 mm.

of *D. poncei* is faint to obsolete whereas in males of *D. bicolor sp. n.* is strong. The legs of *D. poncei* do not contrast in color with the rest of the body as in *D. bicolor sp. n.* It can be distinguished from *D. gertschi* by its higher pectinal tooth count in males (17 to 19 vs 13 to 15 in *D. gertschi*). The frontal notch of the carapace in *D. gertschi* is “V” shaped whereas in *D. bicolor sp. n.* is “U” shaped. Surfaces on the pedipalp patella of *D. gertschi* are finely granular and punctate (this condition is present in the species of the genus *Didymocentrus*, some species of the genus *Bioculus* and has been reported only in another three species in the genus *Diplocentrus*, Sissom & Walker, 1992; Armas & Martín-Friás, 2004; Armas *et al.*, 2004; Santibáñez-López, pers. obs.) whereas in *D. bicolor sp. n.* they are shagreened to granular, without punctuation; and *D. gertschi* (adults: 45 to 52 mm) is smaller than *D. bicolor sp. n.* It can be distinguished from *D. zacatecanus* by a higher telotarsal formula (7/8 7/8 on the last two legs, whereas on *D. zacatecanus* they are 6/7 7/7), and a higher pectinal tooth count in males (17 to 19 with a mode of 18, whereas on *D. zacatecanus*, it is 12 to 16 with a mode of 13), and in females (15 to 16, with a mode of 14 whereas on *D. zacatecanus*, it is 8–14 with a mode of 11 to 12). The frontal notch of the carapace in *D. zacatecanus* is “V” shaped whereas in *D. bicolor sp. n.* is “U” shaped.

Description of the holotype male (Figs. 1–2): **Prosoma.** Carapace reddish brown to dark red, shagreened to sparsely and minutely granular. Frontal notch of carapace “U” shaped, deep, slightly granular and setose; three pairs of lateral eyes subequal in size (Fig. 3). Carapace smooth to minutely granular.

Mesosoma. Tergites reddish brown with fusco-piceus pattern (Darker in female). Tergites I–VI surface shagreened to weakly granular towards anterior margin. Tergite VII shagreened towards the anterior margin and minutely granular towards the posterior margin. Tergite VII presents weak lateral carinae. Pectinal tooth count 17–16 (female with 14–15).

Metasoma. Dark red to reddish brown. Ventral submedian carinae: weak, slightly crenulate on segments I–II; weak to vestigial on segments III–IV. Ventral lateral carinae: strong, smooth and pale on segments I–II; strong, granular and darker on segments III–IV. Lateral inframedian carinae: with a few big conical granules on segments I–II; weak to vestigial on segments III–IV. Lateral supramedian carinae: strong, smooth on segment I; moderate, granular on segments II–IV. Dorsolateral carinae: strong, crenulate on segments I–IV. Segment V slightly longer than pedipalp femur or patella (ratio: 0.97); ventral median and ventral lateral carinae strongly granular with large

TERMS OF USE

This pdf is provided by Magnolia Press for private/research use.
Commercial sale or deposition in a public library or website is prohibited.


subconical granules; ventral transverse carinae with 5 subconical granules and 5 large dark granules on the posterior end of the segment; dorsolateral carinae weak, smooth to slightly granular (Carinae stronger in female and more granular).

Telson. Reddish brown to dark red (Paler in female), smooth with scattered granules at the base of the vesicle; densely setose ventrally (length/width, ratio 0.29).

Pedipalp. Reddish brown to dark red, with carinae darker. Orthobothriotaxic type "C" pattern typical for the genus (Francke, 1977). Femur wider than deep (ratio 0.93) (Fig. 4). Dorsal internal carina strong, granular, becoming obsolete towards posteriorly. Ventral external carina only present basally, weakly granular. Dorsal surface flat with few, small, sparse granules on median portion. Ventral surface flat, shagreened. Internal surface granular with large dark granules.



FIGURES 3-7. *Diplocentrus bicolor* sp. n. 3. Carapace of holotype male. 4. Pedipalp femur, dorsal view. 5. Pedipalp patella, external view. 6. Pedipalp chela, holotype male, dorsal external view. 7. Pedipalp chela, paratype female, dorsal external view. Scale bars= 3 mm.



FIGURES 8–10. Paratype male hemispermophore. 8. Ventral view. 9. Dorsal view. Scale bars= 3 mm. 10. Detail of the capsular region. Scale bar= 1 mm.

Patella (Fig. 5): Dorsal internal carina weak to obsolete, with few large granules, one larger than the others. Dorsal medial carina strong, smooth. Dorsal external carina weak to moderate, smooth. Ventral external carina weak, smooth. Ventral medial carina weak to vestigial, smooth. Ventral internal carina strong, granular. Dorsal, external and ventral faces shagreened; internal surface granular.

Chela (Fig 6): Dorsal marginal carina strong, granular. Dorsal secondary carina weak, smooth. Digital carina strong, smooth. External secondary carina weak to moderate, smooth. Ventral external carina originating at the base of the fixed finger and becoming obsolete basally towards the middle portion. Ventral medial carina strong, smooth. Ventral internal carina originating at the base of the fixed finger, weak, smooth. Fixed finger: dorsal surface basally smooth, densely setose, internal surface moderately concave and external surface slightly convex.

Hemispermophore: 6.9 mm in total length, lamellate, distal lamella 4.4 mm long. Capsular region 1.2 mm wide. Opercular “hook” narrow, without spines (Figs. 8–10).

Intraspecific variation. *Diplocentrus bicolor* sp. n. shows sexual dimorphism: females are darker than males, and the pedipalp chela carinae are more strongly developed in males than in females. Pectinal tooth count variation is as follow: in males (n=29): 6 combs with 14, 2 with 15, 13 with 16, 13 with 17, 14 with 18, 8 with 19 and 2 with 20 teeth; in females (n=10): 1 comb with 12, 7 with 14, 7 with 15 and 5 with 16 teeth.

Legs. Pale brown to yellow, smooth .Telotarsal formula: 6/7 6/7: 6/7 7/8: 7/8 7/8: 7/8 8/8.

TERMS OF USE

This pdf is provided by Magnolia Press for private/research use.
Commercial sale or deposition in a public library or website is prohibited.

TABLE 1. Measurements in millimeters of the holotype male and the paratype female two additional adult males and two adult females of *Diplocentrus bicolor* sp. n.

	Holotype	Paratype	male	female	male	female
Total length	75.3	74.7	59.2	65.4	58	62.8
Carapace length	8.1	10	7.5	9.1	7.9	8.6
Mesosoma length	29.8	23.9	17.4	21.9	17.3	21.9
Metasoma length	43.3	41.7	34.3	34.4	32.8	32.3
Telson length	8.4	8.9	7	8.1	6.7	7.6
Vesicle length	7	7.4	6	70	5.7	64
Vesicle width	3.5	3.8	2.8	36	3.3	3.8
Femur length	9.5	9.4	7.5	8.3	7.8	7.4
Width	3	3.9	3.4	3.5	3	3.3
Depth	2.8	2.7	2.3	2.2	2.2	2.5
Patella length	9	8.2	7.5	7.8	7	7.7
Width	3.5	4	3	36	3	3.1
Chela length	17.7	18	16.5	20.4	16.8	16.2
Width	4.6	6.7	4	5.1	4.8	4.6
Depth	6.7	8.9	7	8.4	7.6	7.6
Movable finger length	11.4	10.8	7.4	8	9.6	9.1
Fixed finger length	8.7	8.6	9.5	10	7.8	6.9
Chelicera length	2.9	3.4	3	3.2	3	3
Width	1.9	2.4	1.9	2	2	2.1
Movable finger length	1.7	2.5	1.7	1.9	2	1.8
Fixed finger length	1.4	1.5	1.5	1.4	15	1.3
Pectinal tooth count	18-16	15-14	17-16	16-16	18-19	15-14

Telotarsal spiniform setae count variation as is show in table2 (n=39):

TABLE 2. Telotarsal spiniform setae count:

Leg #/# of Setae	3	4	5	6	7	8	9	Missing
1 prolateral	1		42	31				1
1 retrolateral			1	34	40	2		1
2 prolateral			4	58	14			2
2 retrolateral				5	50	21		2
3 prolateral			1	10	63	4		
3 retrolateral	1			1	19	55	2	
4 prolateral				1	5	59	9	4
4 retrolateral		1	15			50	8	4

Therefore, the modal formula for the species is 5/7:6/7:7/8:7/8.

TERMS OF USE

This pdf is provided by Magnolia Press for private/research use.
Commercial sale or deposition in a public library or website is prohibited.



FIGURE 11. Known distribution of *Diplocentrus* species in western México. Legend: Black circle= *D. zacatecanus*, black cross = *D. gertchi*, black square = *D. poncei*, black triangule = *D. bicolor* sp. n.

Acknowledgements

We would like to thank Dr. O. Francke and Dr. José Luis Navarrete for the review of the first draft of the manuscript. We also thank O. Francke, J. Ponce, E. González, A. Jaimes, G. Francke, V. Capovilla and M. Córdova for collecting all the know specimens. To A. Valdez, J. Cruz and D. Barrales for their help with the photographs. The senior author would like to thank the program of Practicas Profesionales of the Universidad de Guadalajara and the junior author would like to acknowledge CONACYT and the Graduate Program in Biological Sciences of the Universidad Nacional Autónoma de México for financial support. Specimen collections were done under Scientific permit FAUT-0175 from Semarnat to O. Francke. Financial support was also partially provided by U.S. National Science Foundation DEB 0413453 grant to Dr. L. Prendini and by scorpion related research projects financed by Instituto Bioclon to Dr. O. Francke.

References

- Armas, L. & E. Martín-Frías. (2004) Nueva especie de *Diplocentrus* Peters, 1861 (Scorpionidae: Diplocentrinae) de Veracruz, México. *Revista Ibérica de Aracnología*, 9, 103–106.
- Armas, L., E. Martín-Frías & P. Berea. (2004) Nuevo *Diplocentrus* Peters, 1861 (Scorpionidae: Diplocentrinae) del Estado de Veracruz, México. *Revista Ibérica de Aracnología*, 10, 275–280.
- Coddington, J.A., G. Giribet, M.S. Harvey, L. Prendini & D.E. Walter. (2004) Arachnida. pp. 296–318. In: Cracraft, J. and Donoghue, M. (Eds.) Assembling the Tree of Life. Oxford University Press, Oxford. England.
- Francke, O.F. (1977) Scorpions of the genus *Diplocentrus* from Oaxaca, Mexico (Scorpionida, Diplocentridae). *Journal of Arachnology*, 4, 145–200.
- Francke, O.F. (2007) Alacranes (Arachnida: Scorpiones) de Frontera Corozal, en la selva lacandona, Chiapas, México con la descripción de una nueva especie de *Diplocentrus* (Diplocentridae). *Revista Mexicana de Biodiversidad*, 78, 69–77.
- Francke, O.F. & Ponce-Saavedra, J. (2005) A new species of *Diplocentrus* (Arachnida: Scorpiones) from Michoacan, Mexico. *Revista Mexicana de Biodiversidad*, 76, 49–53.

TERMS OF USE

This pdf is provided by Magnolia Press for private/research use.
Commercial sale or deposition in a public library or website is prohibited.

- Francke, O.F. & Quijano-Ravell, A. (2009) Una especie nueva de *Diplocentrus* (Scorpiones: Diplocentridae) del estado de Michoacán, México. *Revista Mexicana de Biodiversidad*, 80, 659–663.
- Ponce-Saavedra J., O.F. Francke & F. Quijano-Ravell. (2009) Nuevos registros y distribución actualizada de *Diplocentrus zacatecanus* Hoffmann, (1931) (Scorpiones: Diplocentridae). *Entomología Mexicana*, 8, 57–60
- Prendini, L. (2003) Revision of the genus *Lisposoma* Lawrence, 1928 (Scorpiones: Bothriuridae). *Insect Systematics and Evolution*, 34, 241–264.
- Prendini, L. & Wheeler, W. (2005) Scorpion higher phylogeny and classification, taxonomic anarchy, and standards for peer review in online publishing. *Cladistics*, 21, 446–494.
- Santibáñez-López, C.E. & Francke, O.F. (2008) A new species of *Diplocentrus* (Arachnida: Scorpiones) from Oaxaca, Mexico. *Zootaxa*, 1742, 53–60.
- Santibáñez-López, C.E. & Sissom, W.D. (2010) A new species of the *Vaejovis eusthenura* group in Oaxaca, Mexico (Scorpiones: Vaejovidae). *Zootaxa*, 2493, 49–58.
- Santibáñez-López, C.E., Francke, O.F. & Córdova-Athanasiadis, M. (2011) The genus *Diplocentrus* Peters (Scorpiones: Diplocentridae) in Morelos, Mexico. *Revista Ibérica de Aracnología*, 19, 3–13.
- Sissom, W.D. & Fet, V. (2000) Family Diplocentridae. pp. 329–354. In: Fet, V., Sissom, W.D., Lowe, G. & Braunwalder, M.E., *Catalog of the Scorpions of the World (1758–1998)*. The New York Entomological Society. New York, USA.
- Sissom, W.D. & Walker, A.L. (1992) A new species of *Diplocentrus* from Western Mexico. *Southwestern Naturalist*, 37(2), 126–131.
- Stahnke, H.L. (1970) Scorpion nomenclature and mensuration. *Entomological News*, 81, 297–316.
- Vachon, M. (1974) Étude des caractères utilisés pour classer les familles et les genres de Scorpions (Arachnides). 1. La trichobothriataxe en Arachnologie, sigles trichobothriaux et types de trichobothriataxe chez les scorpions. *Bulletin du Muséum National d'Histoire Naturelle*, 3, 857–958.
- Volschenk, E. S. (2005) A new technique for examining surface morphosculpture of scorpions. *Journal of Arachnology*, 33, 820–825.